



Designation: E1242 – 23

## Standard Practice for Using Octanol-Water Partition Coefficient to Estimate Median Lethal Concentrations for Fish Due to Narcosis<sup>1</sup>

This standard is issued under the fixed designation E1242; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon ( $\epsilon$ ) indicates an editorial change since the last revision or reapproval.

### 1. Scope

1.1 This practice covers a procedure for estimating the fathead minnow (*Pimephales promelas*) 96-h LC<sub>50</sub> of nonreactive (that is, covalently bonded without unsaturated residues) and nonelectrolytic (that is, require vigorous reagents to facilitate substitution, addition, replacement reactions and are non-ionic, non-dissociating in aqueous solutions) organic chemicals acting solely by narcosis, also referred to as Meyer-Overton toxicity relationship.<sup>2</sup>

1.2 This procedure is accurate for organic chemicals that are toxic due to narcosis and are non-reactive and non-electrolytic. Examples of appropriate chemicals are: alcohols, ketones, ethers, simple halogenated aliphatics, aromatics, and aliphatic substituted aromatics. It is not appropriate for chemicals whose structures include a potential toxiphore (that structural component of a chemical molecule that has been identified to show mammalian toxicity, for example CN is known to be responsible for inactivation of enzymes, NO<sub>2</sub> for decoupling of oxidative phosphorylation, both leading to mammalian toxicity). Examples of chemicals inappropriate for this practice are: carbamates, organophosphates, phenols, beta-gamma unsaturated alcohols, electrophiles, and quaternary ammonium salts.

1.3 *This international standard was developed in accordance with internationally recognized principles on standardization established in the Decision on Principles for the Development of International Standards, Guides and Recommendations issued by the World Trade Organization Technical Barriers to Trade (TBT) Committee.*

<sup>1</sup> This practice is under the jurisdiction of ASTM Committee E50 on Environmental Assessment, Risk Management and Corrective Action and is the direct responsibility of Subcommittee E50.47 on Biological Effects and Environmental Fate.

Current edition approved Jan. 1, 2023. Published February 2023. Originally approved in 1988. Last previous edition approved in 2014 as E1242 – 97(2014). DOI: 10.1520/E1242-23.

<sup>2</sup> Lipnick, Robert L., “Validation and Extension of Fish Toxicity QSARs and Interspecies Comparisons for Certain Classes of Organic Chemicals,” *QSAR in Toxicology and Xenobiochemistry*, Elsevier, 1985.

### 2. Referenced Documents

#### 2.1 ASTM Standards:<sup>3</sup>

E729 Guide for Conducting Acute Toxicity Tests on Test Materials with Fishes, Macroinvertebrates, and Amphibians

E943 Terminology Relating to Biological Effects and Environmental Fate (Withdrawn 2023)<sup>4</sup>

E1023 Guide for Assessing the Hazard of a Material to Aquatic Organisms and Their Uses

E1147 Test Method for Partition Coefficient (N-Octanol/Water) Estimation by Liquid Chromatography (Withdrawn 2013)<sup>4</sup>

### 3. Terminology

#### 3.1 Definitions:

3.1.1 *acute toxicity*—where an adverse effect such as mortality is measured after organisms exposed to a compound for a relatively short period, usually not constituting a substantial portion of their life span.

3.1.2 *narcosis*, *n*—a reversible state of stupor, unconsciousness, or arrested activity produced by the influence of chemicals on critical sites within membranes or by disrupting the normal functioning of certain proteins by means of nonspecific binding of organic chemical(s) to hydrophobic sites. Death results if exposure is not terminated after a length of time which varies with concentration.

3.1.3 *octanol-water partition coefficient* ( $K_{ow}$ ), *n*—referred to as *P* in some literature.

3.1.4 *toxiphore*, *n*—a chemical structure substituent group that when present gives rise to an adverse effect in exposed organisms.

<sup>3</sup> For referenced ASTM standards, visit the ASTM website, [www.astm.org](http://www.astm.org), or contact ASTM Customer Service at [service@astm.org](mailto:service@astm.org). For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.

<sup>4</sup> The last approved version of this historical standard is referenced on [www.astm.org](http://www.astm.org).