



Designation: E1482 – 23

Standard Practice for Use of Gel Filtration Columns for Cytotoxicity Reduction and Neutralization^{1,2}

This standard is issued under the fixed designation E1482; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This practice is intended to be used to reduce the cytotoxic level of the virus-test product mixture prior to assaying for viral infectivity. It is used in conjunction with evaluations of the virucidal efficacy of disinfectant solutions, wipes, trigger sprays, or pressurized disinfectant spray products intended for use on inanimate, nonporous environmental surfaces. This practice may also be used in the evaluation of hygienic handwashes/handrubs, or for other special applications. The practice may be employed with all viruses and host systems.

NOTE 1—Gel filtration columns may impact virus titer and their use should be taken into consideration when selected for use.

1.2 This practice should be performed only by persons trained in virology techniques.

1.3 This practice utilizes gel filtration technology. The effectiveness of the practice is dependent on the ratio of gel bed volume to sample size and uniformity in the preparation of columns as well as the conditions of centrifugation. The effectiveness of this practice is maximized by investigator practice and experience with gel filtration techniques.

1.4 This practice will aid in the reduction, but not necessarily elimination, of test product toxicity while preserving the titer of the input virus.

1.5 *Units*—The values stated in SI units are to be regarded as standard. No other units of measurement are included in this standard.

1.6 *This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety, health, and environmental practices and determine the applicability of regulatory limitations prior to use.*

¹ This practice is under the jurisdiction of ASTM Committee E35 on Pesticides, Antimicrobials, and Alternative Control Agents and is the direct responsibility of Subcommittee E35.15 on Antimicrobial Agents.

Current edition approved April 1, 2023. Published April 2023. Originally approved in 1992. Last previous edition approved in 2017 as E1482 – 12 (2017). DOI: 10.1520/E1482-23.

² The title was formerly Standard Test Method for Neutralization of Virucidal Agents in Virucidal Efficacy Evaluations.

1.7 *This international standard was developed in accordance with internationally recognized principles on standardization established in the Decision on Principles for the Development of International Standards, Guides and Recommendations issued by the World Trade Organization Technical Barriers to Trade (TBT) Committee.*

2. Referenced Documents

2.1 *ASTM Standards*:³

E1052 Practice to Assess the Activity of Microbicides against Viruses in Suspension

E1053 Practice to Assess Virucidal Activity of Chemicals Intended for Disinfection of Inanimate, Nonporous Environmental Surfaces

E2756 Terminology Relating to Antimicrobial and Antiviral Agents

3. Terminology

3.1 *Definitions*:

3.1.1 For definitions of terms used in this guide, refer to Terminology E2756.

4. Summary of Test Practice

4.1 After the exposure of a virus to a test product, the virus-product suspension is added to a column of Sephadex⁴ LH-60, Sephadex⁴ LH-20, or other appropriate gel filtration medium. The column (encased within a sterile centrifuge tube in order to capture the filtrate) is placed in a centrifuge and centrifuged to separate the virus from the test product by gel filtration. Alternatively, samples may be hand-plunged using a syringe plunger. The filtrate (the column flow-through which contains the virus) is assayed in the appropriate host system. The untreated virus control suspension is gel-column filtered,

³ For referenced ASTM standards, visit the ASTM website, www.astm.org, or contact ASTM Customer Service at service@astm.org. For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.

⁴ Sephadex is a registered trademark of Amersham Biosciences. The sole source of supply of the apparatus known to the committee at this time is Amersham Biosciences. If you are aware of alternative suppliers, please provide this information to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee,¹ which you may attend.